

# MODEL EQUATIONS OF DIFFUSIVE MICROSTRETCH THERMOELASTICITY WITH MICROTERTURES AND MICROCONCENTRATIONS

A. Chirilă<sup>1</sup> and M. Marin<sup>1</sup>

<sup>1</sup>Department of Mathematics and Computer Science, Transilvania University of Braşov, Braşov, Romania

e-mail: adina.chirila@unitbv.ro

## 1. Motivation

This paper is dealing with the linear theory of microstretch thermoelasticity, a theory that is part of the mechanics of generalized continua and which accurately describes the behaviour of materials with microstructure. The concept of generalized continua was introduced by the Cosserat brothers in 1909, who considered a micropolar continuum to be a collection of interconnected particles in the form of small rigid bodies that have three additional rotational degrees of freedom besides the three translational degrees of freedom from classical continuum mechanics. Cosserat thermoelasticity was generalized to microstretch thermoelasticity by Eringen in 1990 [6] by including the effect of axial stretch during the rotation of molecules.

In this paper, we prove the well-posedness of the mathematical model introduced in [3] for the anisotropic case. The idea is to verify the assumptions from the Lumer-Phillips corollary to the Hille-Yosida theorem. As in [2], the theory we derived is closer to the realistic constitutive structure of solids since it describes thermal-diffusion interactions at the macroscopic and microscopic levels. Given the increasing interest in nanomaterials, it is important to take into consideration both the microtemperatures and the microconcentrations of the nanoparticles. The concept of microconcentrations is a novel one, introduced for the first time in the theory of thermoelasticity in [2] and for the first time in mechanics of generalized continua in [3].

## 2. Basic equations

The equations of diffusive microstretch thermoelasticity with microtemperatures and microconcentrations [3]

$$(1) \quad \begin{aligned} t_{ji,j} + \rho f_i &= \rho \ddot{u}_i & \rho T_0 \dot{S} &= q_{i,i} + \rho s \\ h_{k,k} + g + \rho l &= J \ddot{\varphi} & \rho \dot{\varepsilon}_i &= q_{ji,j} + q_i - Q_i + \rho G_i & \dot{C} &= \eta_{i,i} \\ m_{ji,j} + \varepsilon_{irs} t_{rs} + \rho g_i &= I_{ij} \ddot{\varphi}_j & \rho \dot{\omega}_i &= \eta_{ji,j} + \eta_i - \tilde{\sigma}_i \end{aligned}$$

In the equations above,  $u_i$  is the displacement vector field,  $\varphi$  is the microdilatation function,  $\varphi_i$  is the microrotation vector,  $t_{ij}$  is the stress tensor,  $\rho$  is the reference mass density,  $f_i$  is the body force,  $h_j$  is the microstretch vector,  $g$  is the internal body force,  $l$  is the external microstretch body load,  $m_{ij}$  is the couple stress tensor,  $g_i$  is the body couple density,  $C$  is the concentration,  $\eta_i$  is the flux vector of mass diffusion,  $S$  is the microentropy. According to [3], the constitutive equations of the mathematical model for the anisotropic case are

$$(2) \quad \begin{aligned} t_{ij} &= A_{ijrs} e_{rs} + B_{ijrs} \kappa_{rs} + D_{ij} \varphi + F_{ijk} \zeta_k + L_{ijk} T_k - a_{ij} \theta + d_{ij} C \\ m_{ij} &= B_{rsij} e_{rs} + C_{ijrs} \kappa_{rs} + E_{ij} \varphi + G_{ijk} \zeta_k + M_{ijk} T_k - b_{ij} \theta + f_{ij} C \\ h_i &= F_{rsi} e_{rs} + G_{rsi} \kappa_{rs} + A_{ij} \zeta_j + B_i \varphi - N_{ij} T_j - d_i \theta + \tilde{f}_i C \\ g &= -D_{ij} e_{ij} - E_{ij} \kappa_{ij} - B_i \zeta_i - \xi \varphi - R_i T_i + F \theta - \tilde{g}_1 C \\ \rho S &= a_{ij} e_{ij} + b_{ij} \kappa_{ij} + d_i \zeta_i + F \varphi + b_i T_i + a \theta + \varpi C \\ \rho \varepsilon_i &= L_{rsi} e_{rs} + M_{rsi} \kappa_{rs} - N_{ji} \zeta_j + R_i \varphi - B_{ij} T_j - b_i \theta - R_{ij} C_j \\ P &= d_{ij} e_{ij} + f_{ij} \kappa_{ij} + \tilde{f}_i \zeta_i + \tilde{g}_1 \varphi - \varpi \theta + \varrho C & \rho \omega_i &= -C_{ij} C_j - R_{ji} T_j \end{aligned}$$

Here,  $T_0$  is the absolute temperature in the reference configuration,  $q_i$  is the heat flux vector,  $s$  is the heat supply per unit mass,  $\varepsilon_i$  is the first moment of energy vector,  $q_{ij}$  is the first heat flux moment tensor,  $Q_i$  is the microheat

flux average,  $G_i$  is the first heat supply moment vector,  $\eta_{ij}$  is the first mass diffusion flux moment tensor,  $\tilde{\sigma}_i$  is the micromass diffusion flux average,  $P$  is the particle chemical potential,  $T$  is the absolute temperature,  $\theta = T - T_0$ ,  $T_i$  are the microtemperatures,  $C_i$  are the microconcentrations and  $\varepsilon_{ijk}$  is the alternating symbol.

### 3. Well-posedness

We assume null boundary conditions. We introduce the notations  $\dot{u}_i = v_i$ ,  $\dot{\varphi}_i = \psi_i$  and  $\dot{\varphi} = \psi$ . Let  $\mathcal{H} = \{(u_i, v_i, \varphi, \psi, \varphi_i, \psi_i, \theta, T_i, P, C_i) : u_i, \varphi_i \in \mathbb{W}_0^{1,2}(\Omega), v_i, \psi_i, T_i, C_i \in \mathbb{L}^2(\Omega), \psi, \theta, P \in L^2(\Omega), \varphi \in W_0^{1,2}(\Omega)\}$  where  $W_0^{1,2}(\Omega)$ ,  $L^2(\Omega)$  are the familiar Sobolev spaces and  $\mathbb{W}_0^{1,2}(\Omega) = [W_0^{1,2}(\Omega)]^3$ ,  $\mathbb{L}^2(\Omega) = [L^2(\Omega)]^3$ . The boundary initial value problem can be transformed into the following equation in the Hilbert space  $\mathcal{H}$

$$(3) \quad \frac{d\mathcal{U}}{dt} = \mathcal{A}\mathcal{U}(t) + \mathcal{F}(t) \quad \mathcal{U}(0) = \mathcal{U}_0$$

where  $\mathcal{U} = (u_i, v_i, \varphi, \psi, \varphi_i, \psi_i, \theta, P, T_i, C_i)$ ,  $\mathcal{U}_0 = (u_i^0, v_i^0, \varphi^0, \psi^0, \varphi_i^0, \psi_i^0, \theta^0, P^0, T_i^0, C_i^0)$  is the vector of initial conditions and  $\mathcal{A}$  is a certain matrix operator on  $\mathcal{H}$ .

**Lemma 3.1** *In the case of diffusive microstretch thermoelasticity with microtemperatures and microconcentrations and for every  $\mathcal{U} \in \mathcal{D}(\mathcal{A})$ , the operator  $\mathcal{A}$  satisfies the inequality*

$$(4) \quad \langle \mathcal{A}\mathcal{U}, \mathcal{U} \rangle \leq 0$$

in a suitably introduced inner product in  $\mathcal{H}$ .

**Lemma 3.2** *In the case of diffusive microstretch thermoelasticity with microtemperatures and microconcentrations, and for  $Id$  the identity operator in  $\mathcal{H}$ , the operator  $\mathcal{A}$  has the property that*

$$(5) \quad \text{Range}(Id - \mathcal{A}) = \mathcal{H}$$

**Theorem 3.1** *The operator  $\mathcal{A}$  generates a semigroup of contractions in  $\mathcal{H}$ .*

This result proves that in the motion following any sufficiently small change in the external system, the solution of the initial boundary value problem is everywhere arbitrary small in magnitude. Now that we proved that the mathematical model is well-posed by means of the semigroup of linear operator theory, the asymptotic behaviour of solutions and the effect of a concentrated heat source can be studied.

### References

- [1] M. Aouadi. Some theorems in the isotropic theory of microstretch thermoelasticity with microtemperatures. *Journal of Thermal Stresses*, 31(7):649-662, 2008.
- [2] M. Aouadi, M. Ciarletta, and V. Tibullo. A thermoelastic diffusion theory with microtemperatures and microconcentrations. *Journal of Thermal Stresses*, 40(4):486-501, 2017.
- [3] A. Chirilă and M. Marin. Diffusion in microstretch thermoelasticity with microtemperatures and microconcentrations. in *Springer book's series Studies in Systems, Decision and Control, Contributed Volume: Advanced Models and Theories in Social Systems*, accepted, 2018.
- [4] A. Chirilă and M. Marin. The theory of generalized thermoelasticity with fractional order strain for dipolar materials with double porosity. *Journal of Materials Science*, 53(5):3470-3482, 2018.
- [5] C. M. Dafermos. Contraction semigroups and trends to equilibrium in continuum mechanics. in P. German and B. Nayroles (eds.) *Applications of methods of functional analysis to problems in mechanics*, Springer Lecture Notes in Math., 503:295-306, Springer-Verlag, Berlin, 1976.
- [6] A. C. Eringen. Theory of thermo-microstretch elastic solids. *International Journal of Engineering Science*, 28(12):1291-1301, 1990.
- [7] M. Marin, I. Abbas, and R. Kumar. Relaxed Saint-Venant principle for thermoelastic micropolar diffusion. *Structural Engineering and Mechanics*, 51(4):651-662, 2014.
- [8] M. Marin and S. Nicaise. Existence and stability results for thermoelastic dipolar bodies with double porosity. *Continuum Mechanics and Thermodynamics*, 28(6):1645-1657, 2016.