743.

ON THE NEWTON-FOURIER IMAGINARY PROBLEM.

[From the Proceedings of the Cambridge Philosophical Society, vol. III. (1880), pp. 231, 232.]

THE Newtonian process of approximation to the root of a numerical equation f(u) = 0, consists in deriving from an assumed approximate root ξ a new value $\xi_1 = \xi - \frac{f(\xi)}{f'(\xi)}$, which should be a closer approximation to the root sought for: taking the coefficients of f(u) to be real, and also the root sought for, and the assumed value ξ , to be each of them real, Fourier investigated the conditions under which ξ_1 is in fact a closer approximation. But the question may be looked at in a more general manner: ξ may be any real or imaginary value, and we have to inquire in what cases the series of derived values

$$\xi_1 = \xi - \frac{f(\xi)}{f'(\xi)}, \quad \xi_2 = \xi_1 - \frac{f(\xi_1)}{f'(\xi_1)}, \dots$$

converge to a root, real or imaginary, of the equation f(u)=0. Representing as usual the imaginary value ξ , =x+iy, by means of the point whose coordinates are x, y, and in like manner ξ_1 , $=x_1+iy_1$, &c., then we have a problem relating to an infinite plane; the roots of the equation are represented by points A, B, C, \ldots ; the value ξ is represented by an arbitrary point P; and from this by a determinate geometrical construction we obtain the point P_1 , and thence in like manner the points P_2 , P_3 ,... which represent the values ξ_1 , ξ_2 , ξ_3 ,... respectively. And the problem is to divide the plane into regions, such that, starting with a point P_1 anywhere in one region, we arrive ultimately at the root A; anywhere in another region we arrive ultimately at the root B; and so on for the several roots of the equation. The division into regions is made without difficulty in the case of a quadric equation; but in the next succeeding case, that of a cubic equation, it is anything but obvious what the division is : and the author had not succeeded in finding it.